arXiv:1508.06871v2 [math.NA] 12 Jun 2016

Discrete Green functions of the SDFEM on Shishkin triangular meshes

Jin Zhang*†

June 14, 2016

Abstract

We present bounds of discrete Green functions in the energy norm for the standard (or modified) streamline diffusion finite element method (SDFEM) on Shishkin triangular meshes.

1 Problem

We consider the singularly perturbed boundary value problem

$$\begin{aligned} -\varepsilon\Delta u + \boldsymbol{b}\cdot\nabla u + cu = f \quad &\text{in} \quad \Omega=(0,1)^2, \\ u = 0 \quad &\text{on} \quad \partial\Omega, \end{aligned} \tag{1.1}$$

where $\varepsilon \ll |\boldsymbol{b}|$ is a small positive parameter, $\boldsymbol{b}=(b_1,b_2)^T$ is a constant vector with $b_1>0, b_2>0$ and $c>0$ is constant. It is also assumed that f is sufficiently smooth. The solution of (1.1) typically has two exponential layers of width $O(\varepsilon\ln(1/\varepsilon))$ at the sides $x=1$ and $y=1$ of Ω.

2 The SDFEM on Shishkin meshes

2.1 Shishkin meshes

When discretizing (1.1), we use *Shishkin* meshes, which are piecewise uniform. See [5, 7, 2] for a detailed discussion of their properties and applications.

First, we define two mesh transition parameters, which are to be used to specify the mesh changes from coarse to fine in $x-$ and $y-$direction,

$$\lambda_x := \min\left\{\frac{1}{2}, \rho\frac{\varepsilon}{\beta_1}\ln N\right\} \quad \text{and} \quad \lambda_y := \min\left\{\frac{1}{2}, \rho\frac{\varepsilon}{\beta_2}\ln N\right\},$$

---

*Email: jinzhangalex@hotmail.com

†School of Mathematical Sciences, Shandong Normal University, Jinan 250014, China



Figure 1: Dissection of Ω and triangulation $\mathcal{T}_N$.



Figure 2: $K^1_{i,j}$ and $K^2_{i,j}$

where β_1 and β_2 are defined as in [3, Assumption 2.1]. For technical reasons, we set $\rho = 2.5$ in our analysis which is the same as ones in [9] and [8]. The domain Ω is dissected into four parts as $\bar{\Omega} = \Omega_s \cup \Omega_x \cup \Omega_y \cup \Omega_{xy}$ (see Figure 1), where

$$\Omega_s := [0, 1-\lambda_x] \times [0, 1-\lambda_y], \qquad \Omega_x := [1-\lambda_x, 1] \times [0, 1-\lambda_y],$$
$$\Omega_y := [0, 1-\lambda_x] \times [1-\lambda_y, 1], \qquad \Omega_{xy} := [1-\lambda_x, 1] \times [1-\lambda_y, 1].$$

Assumption 1. *We assume that $\varepsilon \leqslant N^{-1}$, as is generally the case in practice. Furthermore we assume that $\lambda_x = \rho\varepsilon\beta_1^{-1}\ln N$ and $\lambda_y = \rho\varepsilon\beta_2^{-1}\ln N$ as otherwise N^{-1} is exponentially small compared with ε.*

Next, we define the set of mesh points $\{(x_i, y_j) \in \Omega : i,\, j = 0, \cdots, N\}$

$$x_i = \begin{cases} 2i(1-\lambda_x)/N & \text{for } i = 0, \cdots, N/2, \\ 1 - 2(N-i)\lambda_x/N & \text{for } i = N/2+1, \cdots, N \end{cases}$$

and

$$y_j = \begin{cases} 2j(1-\lambda_y)/N & \text{for } j = 0, \cdots, N/2, \\ 1 - 2(N-j)\lambda_y/N & \text{for } j = N/2+1, \cdots, N. \end{cases}$$

By drawing lines through these mesh points parallel to the x-axis and y-axis the domain Ω is partitioned into rectangles. Each rectangle is divided into two triangles by drawing the diagonal which runs from (x_i, y_{j+1}) to (x_{i+1}, y_j). This yields a triangulation of Ω denoted by $\mathcal{T}_N$(see Fig. 1). The mesh sizes $h_{x,i} := x_{i+1} - x_i$ and $h_{y,j} := y_{j+1} - y_j$ satisfy

$$h_{x,i} = \begin{cases} H_x := \dfrac{1-\lambda_x}{N/2} & \text{for } i = 0, \cdots, N/2-1, \\[2ex] h_x := \dfrac{\lambda_x}{N/2} & \text{for } i = N/2, \cdots, N-1 \end{cases}$$

and

$$h_{y,j} = \begin{cases} H_y := \dfrac{1-\lambda_y}{N/2} & \text{for } j = 0, \cdots, N/2-1, \\ h_y := \dfrac{\lambda_y}{N/2} & \text{for } j = N/2, \cdots, N-1. \end{cases}$$

The mesh sizes $h_{x,i}$ and $h_{y,j}$ satisfy

$$N^{-1} \leqslant H_x, H_y \leqslant 2N^{-1} \quad \text{and} \quad C_1 \varepsilon N^{-1} \ln N \leqslant h_x, h_y \leqslant C_2 \varepsilon N^{-1} \ln N.$$

For convenience, we shall use some notations: $K_{i,j}^1$ for the mesh triangle with vertices (x_i, y_j), (x_{i+1}, y_j) and (x_i, y_{j+1}); $K_{i,j}^2$ for the mesh triangle with vertices (x_i, y_{j+1}), (x_{i+1}, y_j) and (x_{i+1}, y_{j+1}) (see Fig. 2); τ or K for a generic mesh triangle.

2.2 The streamline diffusion finite element method

On the above Shishkin meshes we define a C^0 linear finite element space

$$V^N := \{v^N \in C(\bar{\Omega}) : v^N|_{\partial\Omega} = 0 \text{ and } v^N|_K \in P_1(K), \forall K \in \mathcal{T}_N\}.$$

Now we are in a position to state the standard SDFEM for (1.1) which reads:

$$\begin{cases} \text{Find } u^N \in V^N \text{ such that for all } v^N \in V^N, \\ a_{SD}(u^N, v^N) = (f, v^N) + \displaystyle\sum_{K \subset \Omega} (f, \delta_K \boldsymbol{b} \cdot \nabla v^N)_K, \end{cases} \tag{2.1}$$

where

$$\begin{aligned} a_{SD}(u^N, v^N) =& \varepsilon(\nabla u^N, \nabla v^N) + (\boldsymbol{b} \cdot \nabla u^N, v^N) + (cu^N, v^N) \\ &+ \sum_{K \subset \Omega} (-\varepsilon \Delta u^N + \boldsymbol{b} \cdot \nabla u^N + cu^N, \delta_K \boldsymbol{b} \cdot \nabla v^N)_K. \end{aligned}$$

Note that $\Delta(u^N|_K) = 0$ for $u^N|_K \in P_1(K)$. Following usual practice [7], the parameter $\delta_K := \delta|_K$ is defined as follows

$$\delta := \delta(x, y) = \begin{cases} C^* N^{-1} & \text{if } (x, y) \in \Omega_s, \\ 0 & \text{otherwise,} \end{cases} \tag{2.2}$$

where C^* is referred to [7, Lemma 3.25].

We set

$$b := \sqrt{b_1^2 + b_2^2}, \quad \boldsymbol{\beta} := \begin{pmatrix} b_1 \\ b_2 \end{pmatrix} / b, \quad \boldsymbol{\eta} := \begin{pmatrix} -b_2 \\ b_1 \end{pmatrix} / b \quad \text{and} \quad v_\zeta := \boldsymbol{\zeta}^T \nabla v$$

for any vector $\boldsymbol{\zeta}$ of unit length. By an easy calculation one shows that

$$(\nabla w, \nabla v) = (w_\beta, v_\beta) + (w_\eta, v_\eta).$$

We rewrite (2.1) as

$$\varepsilon(u_\beta^N, v_\beta^N) + \varepsilon(u_\eta^N, v_\eta^N) + (bu_\beta^N + u^N, v^N) + \sum_{K\subset\Omega}(bu_\beta^N + cu^N, \delta_K bv_\beta^N)_K$$
$$= (f, v^N) + \sum_{K\subset\Omega}(f, \delta_K bv_\beta^N)_K.$$

For technical reasons in the later analysis, we increase the crosswind diffusion(see [1, 4, 6]) by replacing $\varepsilon(u_\eta^N, v_\eta^N)$ by $\hat{\varepsilon}(u_\eta^N, v_\eta^N)$ where $\varepsilon \leqslant \hat{\varepsilon} \leqslant N^{-1}$ and $\hat{\varepsilon}$ is constant on each of subdomains including $\Omega_s, \Omega\backslash\Omega_s$. For convenience, we denote $\hat{\varepsilon}|_{\Omega_s}$by $\hat{\varepsilon}_s$. We will consider two cases: (1) $\hat{\varepsilon} = \varepsilon$; (2) $\hat{\varepsilon}$ defined as in [3, pg 463], that is

$$\hat{\varepsilon} = \begin{cases} \tilde{\varepsilon} & \text{if } \boldsymbol{x} \in \Omega_s, \\ \varepsilon & \text{if } \boldsymbol{x} \in \Omega\backslash\Omega_s \end{cases}$$

where $\tilde{\varepsilon} := \max\left(\varepsilon, N^{-3/2}\right)$.

We now state a streamline diffusion method with artificial crosswind diffusion (ACD), also called a modified SDFEM:

$$\begin{cases} \text{Find } u^N \in V^N \text{ such that for all } v^N \in V^N \\ a_{MSD}(u^N, v^N) = (f, v^N + \delta bv_\beta^N), \end{cases}$$

with

$$\begin{aligned} a_{MSD}(u^N, v^N) :=& \varepsilon(u_\beta^N, v_\beta^N) + \hat{\varepsilon}(u_\eta^N, v_\eta^N) + (bu_\beta^N + u^N, v^N) \\ &+ \sum_{K\subset\Omega}(bu_\beta^N + u^N, \delta_K bv_\beta^N)_K. \end{aligned} \tag{2.3}$$

Finally, we define a special energy norm associated with $a_{MSD}(\cdot,\cdot)$:

$$\|v^N\|_{MSD}^2 := \varepsilon\|v_\beta^N\|^2 + \hat{\varepsilon}\|v_\eta^N\|^2 + \|v^N\|^2 + \sum_{K\subset\Omega}\delta_K\|bv_\beta^N\|_K^2, \quad \forall v^N \in V^N.$$

For brevity, we often write $\varepsilon(u_\beta^N, v_\beta^N) + \sum_{K\subset\Omega}(bu_\beta^N, \delta_K\, bv_\beta^N)_K$ as $(\varepsilon + b^2\delta)(u_\beta^N, v_\beta^N)$.

3 The discrete Green function

Let $\boldsymbol{x}^*$ be a mesh node in Ω. The discrete Green's function $G \in V^N$ associated with $\boldsymbol{x}^*$ is defined by

$$a_{MSD}(v^N, G) = v^N(\boldsymbol{x}^*) \quad \forall v^N \in V^N. \tag{3.1}$$

The weight function ω is defined by

$$\omega(\boldsymbol{x}) := g\left(\frac{(\boldsymbol{x}-\boldsymbol{x}^*)\cdot\boldsymbol{\beta}}{\sigma_\beta}\right) g\left(\frac{(\boldsymbol{x}-\boldsymbol{x}^*)\cdot\boldsymbol{\eta}}{\sigma_\eta}\right) g\left(-\frac{(\boldsymbol{x}-\boldsymbol{x}^*)\cdot\boldsymbol{\eta}}{\sigma_\eta}\right)$$

where

$$g(r) = \frac{2}{1+e^r} \quad \text{for } r \in (-\infty, \infty).$$

Now, we are to derive a global estimate on G in the weighted energy norm

$$\begin{aligned} |||G|||_\omega^2 :=& \varepsilon\|\omega^{-1/2}G_\beta\|^2 + \hat{\varepsilon}\|\omega^{-1/2}G_\eta\|^2 + \frac{b}{2}\|(\omega^{-1})_\beta^{1/2}G\|^2 \\ &+ c\|\omega^{-1/2}G\|^2 + \sum_K b^2\delta_K\|\omega^{-1/2}G_\beta\|_K^2. \end{aligned} \tag{3.2}$$

From (2.3) and (3.2), we have

$$\begin{aligned} |||G|||_\omega^2 =& a_{MSD}(\omega^{-1}G, G) - \varepsilon((\omega^{-1})_\beta G, G_\beta) - \hat{\varepsilon}((\omega^{-1})_\eta G, G_\eta) \\ &- \sum_K (b(\omega^{-1})_\beta G + c\omega^{-1}G, \delta_K\, bG_\beta)_K. \end{aligned} \tag{3.3}$$

Considering (3.1) we also have

$$\begin{aligned} a_{MSD}(\omega^{-1}G, G) &= a_{MSD}(E, G) + a_{MSD}((\omega^{-1}G)^I, G) \\ &= a_{MSD}(E, G) + (\omega^{-1}G)(\boldsymbol{x}^*) \end{aligned} \tag{3.4}$$

where $E := \omega^{-1}G - (\omega^{-1}G)^I$.

Lemma 3.1. *If*

$$\sigma_\beta \geqslant k(\varepsilon + \delta_M) \quad \textit{and} \quad \sigma_\eta \geqslant k\hat{\varepsilon}_M^{1/2}, \tag{3.5}$$

where $\delta_M := \max_{\boldsymbol{x}\in\Omega} \delta$ *and* $\hat{\varepsilon}_M := \max_{\boldsymbol{x}\in\Omega} \hat{\varepsilon}$*, then for* $k > 1$ *sufficiently large and independent of* N *and* ε*, we have*

$$a_{MSD}(\omega^{-1}G, G) \geqslant \frac{1}{4}|||G|||_\omega^2.$$

Proof. Hölder inequalities, Cauchy inequalities and [3, Lemma 4.1] give

$$\begin{aligned} &(\varepsilon + b^2\delta)\left|((\omega^{-1})_\beta G, G_\beta)\right| \\ \leqslant& C\varepsilon^{1/2}\sigma_\beta^{-1/2}\|(\omega^{-1})_\beta^{1/2}G\| \cdot \varepsilon^{1/2}\|\omega^{-1/2}G_\beta\| \\ &+ \sum_K Cb\delta_K^{1/2}\sigma_\beta^{-1/2}\|(\omega^{-1})_\beta^{1/2}G\|_K \cdot b\delta_K^{1/2}\|\omega^{-1/2}G_\beta\|_K \\ \leqslant& \frac{1}{2}\varepsilon\|\omega^{-1/2}G_\beta\|^2 + C\varepsilon\sigma_\beta^{-1}\|(\omega^{-1})_\beta^{1/2}G\|^2 + \frac{1}{4}\sum_K b^2\delta_K\|\omega^{-1/2}G_\beta\|_K^2 \\ &+ C\sum_K \delta_K\sigma_\beta^{-1}\|(\omega^{-1})_\beta^{1/2}G\|^2 \\ \leqslant& \frac{1}{2}\varepsilon\|\omega^{-1/2}G_\beta\|^2 + \frac{1}{4}\sum_K b^2\delta_K\|\omega^{-1/2}G_\beta\|_K^2 + C(\varepsilon + \delta)\sigma_\beta^{-1}\|(\omega^{-1})_\beta^{1/2}G\|^2. \end{aligned}$$

Similarly, we have

$$
\begin{aligned}
\hat{\varepsilon}\left|((\omega^{-1})_\eta G, G_\eta)\right| &\leqslant C\hat{\varepsilon}^{1/2}\sigma_\eta^{-1}\cdot\|\omega^{-1/2}G\|\cdot\hat{\varepsilon}^{1/2}\|\omega^{-1/2}G_\eta\| \\
&\leqslant \frac{1}{2}\hat{\varepsilon}\|\omega^{-1/2}G_\eta\|^2 + \frac{1}{2}C\hat{\varepsilon}\sigma_\eta^{-2}\;\; c\|\omega^{-1/2}G\|^2
\end{aligned}
$$

and

$$
\begin{aligned}
\sum_K (c\omega^{-1}G, \delta_K\; bG_\beta)_K &\leqslant \sum_K c^{1/2}\delta_K^{1/2}\;\; c^{1/2}\|\omega^{-1/2}G\|_K\cdot b\delta_K^{1/2}\|\omega^{-1/2}G_\beta\|_K \\
&\leqslant \frac{1}{4}c\|\omega^{-1/2}G\|^2 + \sum_K c\delta_K\cdot b^2\delta_K\|\omega^{-1/2}G_\beta\|_K^2 \\
&\leqslant \frac{1}{4}c\|\omega^{-1/2}G\|^2 + \frac{1}{2}\sum_K b^2\delta_K\|\omega^{-1/2}G_\beta\|_K^2.
\end{aligned}
$$

If (3.5) holds true and k is taken sufficiently large and independent of N and ε, then we have

$$
(\varepsilon+b^2\delta)|((\omega^{-1})_\beta G, G_\beta)| + \varepsilon|((\omega^{-1})_\eta G, G_\eta)| + \sum_K |(c\omega^{-1}G, \delta_K\; bG_\beta)_K| \leqslant \frac{3}{4}|||G|||_\omega^2.
$$

Considering (3.3), we are done.

$\square$

Lemma 3.2. *If* $1 \geqslant \sigma_\beta \geqslant k(\varepsilon+\delta_M)$*, with* $k > 0$ *sufficiently large and independent of* N *and* ε*. Then for each mesh point* $\boldsymbol{x}^* \in \Omega\backslash\Omega_{xy}$*, we have*

$$
\left|(\omega^{-1}G)(\boldsymbol{x}^*)\right| \leqslant \frac{1}{16}|||G|||_\omega^2 + \begin{cases} CN^2\sigma_\beta & \text{if } \boldsymbol{x}^* \in \Omega_s \\ CN\ln N & \text{if } \boldsymbol{x}^* \in \Omega_x \cup \Omega_y \end{cases}
$$

where C *is independent of* N*,* ε *and* $\boldsymbol{x}^*$*.*

Proof. First let $\boldsymbol{x}^* \in \Omega_s$. Let τ^* be the unique triangle that has $\boldsymbol{x}^*$ as its north-east corner. Then

$$
\left|(\omega^{-1}G)(\boldsymbol{x}^*)\right| \leqslant CN\|G\|_{\tau^*} \leqslant CN\max_{\tau^*}\left|(\omega^{-1})_\beta^{-1/2}\right|\cdot\|(\omega^{-1})_\beta^{1/2}G\|_{\tau^*}.
$$

Calculating $(\omega^{-1})_\beta^{-1}(\boldsymbol{x})$ explicitly, we see that

$$
(\omega^{-1})_\beta^{-1}(\boldsymbol{x}) \leqslant C\sigma_\beta \quad \forall \boldsymbol{x} \in \tau^*.
$$

Thus the arithmetic-geometric mean inequality gives

$$
\left|(\omega^{-1}G)(\boldsymbol{x}^*)\right| \leqslant CN^2\sigma_\beta + \frac{1}{16}|||G|||_\omega^2.
$$

Next, let $\boldsymbol{x}^* \in \Omega_x$. (The case $\boldsymbol{x}^* \in \Omega_y$ is similar.) Write $\boldsymbol{x}^* = (x_i, y_j)$. Then

$$
\begin{aligned}
\left|(\omega^{-1}G)(\boldsymbol{x}^*)\right| = |G(\boldsymbol{x}^*)| = \left|\int_{x_i}^1 G_x(t, y_j)\mathrm{d}t\right| &\leqslant CH_y^{-1}\int_{x_i}^1\int_{y_j}^{y_{j+1}} |G_x(t,y)|\,\mathrm{d}y\mathrm{d}t \\
&\leqslant CN(\varepsilon\ln N\cdot N^{-1})^{1/2}\|G_x\|_{\Omega_x} \leqslant CN^{1/2}\ln^{1/2}N|||G||| \\
&\leqslant CN\ln N + \frac{1}{16}|||G|||_\omega^2.
\end{aligned}
$$

Thus, we are done. $\square$

Lemma 3.3. *If*

$$(\varepsilon+b^2\delta)^{1/2}\|\omega^{1/2}E_\beta\|+\hat{\varepsilon}^{1/2}\|\omega^{1/2}E_\eta\|+\delta_s^{-1/2}\|\omega^{1/2}E\|_{\Omega_s} \\ +\varepsilon^{-1/2}\|\omega^{1/2}E\|_{\Omega\setminus\Omega_s}\leqslant Ck^{-1/2}|\!|\!|G|\!|\!|_\omega. \tag{3.6}$$

where $k>1$ sufficiently large and independent of N and ε, then

$$a_{MSD}(E,G)\leqslant\frac{1}{16}|\!|\!|G|\!|\!|_\omega^2$$

where $E=(\omega^{-1}G)^I-\omega^{-1}G$.

Proof. Hölder inequality gives

$$\begin{aligned}|a_{MSD}(E,G)|\leqslant&(\varepsilon+b^2\delta)^{1/2}\|\omega^{1/2}E_\beta\|\cdot(\varepsilon+b^2\delta)^{1/2}\|\omega^{-1/2}G_\beta\|\\&+\hat{\varepsilon}^{1/2}\|\omega^{1/2}E_\eta\|\cdot\hat{\varepsilon}^{1/2}\|\omega^{-1/2}G_\eta\|\\&+C\|\omega^{1/2}E\|\cdot\|\omega^{-1/2}G_\beta\|+\|\omega^{1/2}E\|\cdot\|\omega^{-1/2}G\|.\\&\leqslant\frac{1}{16}|\!|\!|G|\!|\!|_\omega^2,\quad\text{apply (3.6) and take } k \text{ sufficiently large.}\end{aligned}$$

$\square$

Set $h_\tau=\max\{h_{x,\tau},h_{y,\tau}\}$. For $\forall\tau\in\mathcal{T}_N$ we have

$$\|\omega^{1/2}E\|_\tau\leqslant C(\max_\tau\omega)^{1/2}\cdot h_\tau^2\left(\|(\omega^{-1}G)_{\beta\beta}\|_\tau+\|(\omega^{-1}G)_{\beta\eta}\|_\tau+\|(\omega^{-1}G)_{\eta\eta}\|_\tau\right)$$

and

$$\begin{aligned}\|\omega^{1/2}E_\beta\|_\tau\leqslant C(\max_\tau\omega)^{1/2}\cdot h_\tau\left(\|(\omega^{-1}G)_{\beta\beta}\|_\tau+\|(\omega^{-1}G)_{\beta\eta}\|_\tau+\|(\omega^{-1}G)_{\eta\eta}\|_\tau\right),\\ \|\omega^{1/2}E_\eta\|_\tau\leqslant C(\max_\tau\omega)^{1/2}\cdot h_\tau\left(\|(\omega^{-1}G)_{\beta\beta}\|_\tau+\|(\omega^{-1}G)_{\beta\eta}\|_\tau+\|(\omega^{-1}G)_{\eta\eta}\|_\tau\right),\end{aligned}$$

where we have used [3, Corollary 3.1]. Clearly, the following estimate holds

$$\begin{aligned}&\|(\omega^{-1}G)_{\beta\beta}\|_\tau+\|(\omega^{-1}G)_{\beta\eta}\|_\tau+\|(\omega^{-1}G)_{\eta\eta}\|_\tau\\ \leqslant&C\left(\|(\omega^{-1})_{\beta\beta}G\|_\tau+\|(\omega^{-1})_\beta G_\beta\|_\tau\right)\\&+C\left(\|(\omega^{-1})_{\beta\eta}G\|_\tau+\|(\omega^{-1})_\beta G_\eta\|_\tau+\|(\omega^{-1})_\eta G_\beta\|_\tau\right)\\&+C\left(\|(\omega^{-1})_{\eta\eta}G\|_\tau+\|(\omega^{-1})_\eta G_\eta\|_\tau\right).\end{aligned}$$

Set $\alpha_\omega:=(\min_\tau\omega)^{-1/2}$. Lemma 4.1 in [3] yields

$$\begin{aligned}&\|(\omega^{-1})_{\beta\beta}G\|_\tau\leqslant C\alpha_\omega\cdot\sigma_\beta^{-3/2}\cdot\|(\omega^{-1})_\beta^{1/2}G\|_\tau,\\&\|(\omega^{-1})_\beta G_\beta\|_\tau\leqslant C\alpha_\omega\cdot\sigma_\beta^{-1}\cdot\|\omega^{-1/2}G_\beta\|_\tau,\\&\|(\omega^{-1})_{\beta\eta}G\|_\tau\leqslant C\alpha_\omega\cdot\sigma_\beta^{-1/2}\sigma_\eta^{-1}\cdot\|(\omega^{-1})_\beta^{1/2}G\|_\tau,\\&\|(\omega^{-1})_\eta G_\beta\|_\tau\leqslant C\alpha_\omega\cdot\sigma_\eta^{-1}\cdot\|\omega^{-1/2}G_\beta\|_\tau,\\&\|(\omega^{-1})_{\eta\eta}G\|_\tau\leqslant C\alpha_\omega\cdot\sigma_\eta^{-2}\cdot\|\omega^{-1/2}G\|_\tau.\end{aligned}\tag{3.7}$$

Similarly, for $\tau \subset \Omega_s$ we have

$$\|(\omega^{-1})_\eta G_\eta\|_\tau \leqslant C\alpha_\omega \cdot \begin{cases} \sigma_\eta^{-1} N \cdot \|\omega^{-1/2} G\|_\tau & \text{if } \hat{\varepsilon}_s \leqslant N^{-2} \\ \sigma_\eta^{-1}\hat{\varepsilon}_s^{-1/2} \cdot \hat{\varepsilon}_s^{1/2}\|\omega^{-1/2} G_\eta\|_\tau & \text{if } \hat{\varepsilon}_s \geqslant N^{-2} \end{cases} \tag{3.8}$$

and for $\tau \subset \Omega\backslash\Omega_s$

$$\|(\omega^{-1})_\eta G_\eta\|_\tau \leqslant C\alpha_\omega \cdot \varepsilon^{-1/2}\sigma_\eta^{-1} \cdot \varepsilon^{1/2}\|\omega^{-1/2} G_\eta\|_\tau. \tag{3.9}$$

For $\tau \subset \Omega_s$, we have

$$\begin{aligned} \|(\omega^{-1})_\beta G_\eta\|_\tau &\leqslant C\max_\tau(\omega^{-1})_\beta \cdot \|G_\eta\|_\tau \leqslant C\max_\tau(\omega^{-1})_\beta \cdot N \cdot \|G\|_\tau \\ &\leqslant CN \left(\max_\tau(\omega^{-1})_\beta\right)^{1/2} \left(\min_\tau(\omega^{-1})_\beta\right)^{1/2} \cdot \|G\|_\tau \\ &\leqslant C\alpha_\omega \cdot N\sigma_\beta^{-1/2} \cdot \|(\omega^{-1})_\beta^{1/2} G\|_\tau \end{aligned} \tag{3.10}$$

where we have used [3, Lemma 4.1 (vii)]. For $\tau \subset \Omega\backslash\Omega_s$, we have

$$\|(\omega^{-1})_\beta G_\eta\|_\tau \leqslant C\alpha_\omega \cdot \varepsilon^{-1/2}\sigma_\beta^{-1} \cdot \varepsilon^{1/2}\|\omega^{-1/2} G_\eta\|_\tau. \tag{3.11}$$

Set

$$\sigma_\eta(\hat{\varepsilon}) = \begin{cases} \sigma_\eta^{-1} N & \text{if } \hat{\varepsilon}_s \leqslant N^{-2} \\ \sigma_\eta^{-1}\hat{\varepsilon}_s^{-1/2} & \text{if } \hat{\varepsilon}_s \geqslant N^{-2} \end{cases}. \tag{3.12}$$

From (3.7)–(3.11) and [3, Lemma 4.1 (iii)], we have

$$\begin{aligned} &\|\omega^{1/2} E\|_{\Omega_s} + N^{-1}\left(\|\omega^{1/2} E_\beta\|_{\Omega_s} + \|\omega^{1/2} E_\eta\|_{\Omega_s}\right) \\ \leqslant &CN^{-2}\big(\sigma_\beta^{-3/2} + \sigma_\beta^{-1}\delta_s^{-1/2} + \sigma_\beta^{-1/2}\sigma_\eta^{-1} + \sigma_\eta^{-1}\delta_s^{-1/2} + \sigma_\eta^{-2} + \sigma_\eta(\hat{\varepsilon}) + \sigma_\beta^{-1/2}N\big)|||G|||_{\omega,\Omega_s} \\ &\|\omega^{1/2} E\|_{\Omega\backslash\Omega_s} + N^{-1}\left(\|\omega^{1/2} E_\beta\|_{\Omega\backslash\Omega_s} + \|\omega^{1/2} E_\eta\|_{\Omega\backslash\Omega_s}\right) \\ \leqslant &CN^{-2}\big(\sigma_\beta^{-3/2} + \sigma_\beta^{-1}\varepsilon^{-1/2} + \sigma_\beta^{-1/2}\sigma_\eta^{-1} + \sigma_\eta^{-1}\varepsilon^{-1/2} + \sigma_\eta^{-2}\big)|||G|||_{\omega,\Omega\backslash\Omega_s}. \end{aligned} \tag{3.13}$$

To make sure that (3.6) holds, we should have

$$\|\omega^{1/2} E_\beta\|_{\Omega_s} \leqslant Ck^{-1/2}(\varepsilon + \delta_s)^{-1/2}|||G|||_{\omega,\Omega_s}, \tag{3.14}$$

$$\|\omega^{1/2} E_\eta\|_{\Omega_s} \leqslant Ck^{-1/2}\hat{\varepsilon}_s^{-1/2}|||G|||_{\omega,\Omega_s}, \tag{3.15}$$

$$\|\omega^{1/2} E_\beta\|_{\Omega\backslash\Omega_s} + \|\omega^{1/2} E_\eta\|_{\Omega\backslash\Omega_s} \leqslant Ck^{-1/2}\varepsilon^{-1/2}|||G|||_{\omega,\Omega\backslash\Omega_s}, \tag{3.16}$$

$$\|\omega^{1/2} E\|_{\Omega_s} \leqslant Ck^{-1/2}\delta_s^{1/2}|||G|||_{\omega,\Omega_s}, \tag{3.17}$$

and

$$\|\omega^{1/2} E\|_{\Omega\backslash\Omega_s} \leqslant Ck^{-1/2}\varepsilon^{1/2}|||G|||_{\omega,\Omega\backslash\Omega_s}. \tag{3.18}$$

To ensure that (3.14) holds true, we can set

$$\begin{aligned} &\sigma_\beta \geqslant kN^{-1}, \\ &\sigma_\eta \geqslant kN^{-3/4}, \quad \sigma_\eta \geqslant \sigma_\eta^* := \begin{cases} kN^{-1/2} & \text{if } \hat{\varepsilon}_s \leqslant N^{-2} \\ k\hat{\varepsilon}_s^{-1/2}N^{-3/2} & \text{if } \hat{\varepsilon}_s \geqslant N^{-2} \end{cases}. \end{aligned} \tag{3.19}$$

Assume that (3.19) holds true, then (3.14)–(3.17) hold and we have

$$\|\omega^{1/2}E\|_{\Omega\backslash\Omega_s} \leqslant Ck^{-1}\varepsilon^{-1/2}N^{-1}|\!|\!|G|\!|\!|_{\omega}. \tag{3.20}$$

Next, we are to obtain sharper bounds for $\|\omega^{1/2}E\|_{\Omega\backslash\Omega_s}$. Following the techniques of (see [3, Lemma 4.4]), we have

$$(\omega^{1/2}E)(\boldsymbol{x}) = \int_{\boldsymbol{x}}^{\Gamma(\boldsymbol{x})}(\omega^{1/2}E)_\eta \mathrm{d}s \tag{3.21}$$

where $\boldsymbol{x} \in \Omega\backslash\Omega_s$, $\Gamma(\boldsymbol{x}) \in \partial\Omega$ satisfies $(\boldsymbol{x}-\Gamma(\boldsymbol{x}))\cdot\beta = 0$ and the following condition:

$$|\boldsymbol{x}-\Gamma(\boldsymbol{x})| = \min_{y}|\boldsymbol{x}-\boldsymbol{y}|, \text{where } \boldsymbol{y} \in \partial\Omega \text{ and } (\boldsymbol{x}-\boldsymbol{y})\cdot\beta = 0,$$

From (3.21), we have

$$\begin{aligned}\|\omega^{1/2}E\|^2_{\Omega_x\cup\Omega_{xy}} &= \int_{1-\lambda_x}^{1}\int_0^1\left[\int_{\boldsymbol{x}}^{\Gamma(\boldsymbol{x})}(\omega^{1/2}E)_\eta \mathrm{d}s\right]^2 \mathrm{d}y\mathrm{d}x\\ &\leqslant C\lambda_x^2\left\{\|(\omega^{1/2})_\eta E\|^2_{\Omega\backslash\Omega_s} + \|\omega^{1/2}E_\eta\|^2_{\Omega\backslash\Omega_s}\right\}\\ &\leqslant C\varepsilon^2\ln^2 N\left(\sigma_\eta^{-2}\|\omega^{1/2}E\|^2_{\Omega\backslash\Omega_s} + \|\omega^{1/2}E_\eta\|^2_{\Omega\backslash\Omega_s}\right).\end{aligned}$$

Similar argument holds for Ω_y. Thus, we have

$$\|\omega^{1/2}E\|^2_{\Omega\backslash\Omega_s} \leqslant C\varepsilon^2\ln^2 N\left(\sigma_\eta^{-2}\|\omega^{1/2}E\|^2_{\Omega\backslash\Omega_s} + \|\omega^{1/2}E_\eta\|^2_{\Omega\backslash\Omega_s}\right). \tag{3.22}$$

To make sure that (3.18) holds, according to (3.22) the following estimates should hold

$$\begin{aligned}&\varepsilon^2\ln^2 N\sigma_\eta^{-2}\|\omega^{1/2}E\|^2_{\Omega\backslash\Omega_s} \leqslant Ck^{-1}\varepsilon|\!|\!|G|\!|\!|^2_{\omega,\Omega\backslash\Omega_s}\\ &\|\omega^{1/2}E_\eta\|^2_{\Omega\backslash\Omega_s} \leqslant Ck^{-1}\varepsilon^{-1}\ln^{-2}N|\!|\!|G|\!|\!|^2_{\omega,\Omega\backslash\Omega_s}\end{aligned} \tag{3.23}$$

From (3.23) and (3.20), we can set

$$\sigma_\beta \geqslant kN^{-1}\ln N, \quad \sigma_\eta \geqslant kN^{-1}\ln N, \quad \sigma_\eta \geqslant k\varepsilon^{1/4}N^{-1/2}\ln^{1/2}N. \tag{3.24}$$

Assume that (3.19) and (3.24) hold true, substituting (3.20) and (3.23) into the right-hand side of (3.22), we have

$$\|\omega^{1/2}E\|^2_{\Omega\backslash\Omega_s} \leqslant Ck^{-1}\varepsilon|\!|\!|G|\!|\!|^2_{\omega}. \tag{3.25}$$

Thus, we have the following lemma.

Lemma 3.4. *Assume that* σ_β *and* σ_η *satisfy* (3.19) *and* (3.24)*, where* $k > 1$ *sufficiently large and independent of* N *and* ε*, then we have*

$$\begin{aligned}\|\omega^{1/2}E\|_{\Omega_s} &\leqslant Ck^{-1/2}N^{-1/2}|\!|\!|G|\!|\!|_{\omega,\Omega_s},\\ \|\omega^{1/2}E\|_{\Omega\backslash\Omega_s} &\leqslant Ck^{-1/2}\varepsilon^{1/2}|\!|\!|G|\!|\!|_{\omega,\Omega\backslash\Omega_s}\end{aligned}$$

and

$$\begin{aligned}\|\omega^{1/2}E_\beta\|_{\Omega_s}+\|\omega^{1/2}E_\eta\|_{\Omega_s} &\leqslant Ck^{-1/2}N^{1/2}|\!|\!|G|\!|\!|_{\omega,\Omega_s},\\ \|\omega^{1/2}E_\beta\|_{\Omega\backslash\Omega_s}+\|\omega^{1/2}E_\eta\|_{\Omega\backslash\Omega_s} &\leqslant Ck^{-1/2}\varepsilon^{-1/2}\ln^{-1}N|\!|\!|G|\!|\!|_{\omega,\Omega\backslash\Omega_s}.\end{aligned}$$

Theorem 1. *Assume that* σ_β *and* σ_η *satisfy* (3.5)*,* (3.19) *and* (3.24)*, where* k *is chosen so that Lemmas 3.1, 3.2 and 3.3 hold. Then for* $\boldsymbol{x}^* \in \Omega\backslash\Omega_{xy}$*, we have*

$$\|G\|_{MSD} \leqslant \sqrt{8}|\!|\!|G|\!|\!|_\omega \leqslant \begin{cases} CN\sigma_\beta^{1/2} & \textit{if } \boldsymbol{x}^* \in \Omega_s \\ CN^{1/2}\ln^{1/2}N & \textit{if } \boldsymbol{x}^* \in \Omega_x \cup \Omega_y \end{cases}$$

Proof. The readers are referred to [3, Theorem 4.1] for the estimate $\|G\|_{MSD} \leqslant \sqrt{8}|\!|\!|G|\!|\!|_\omega$. Considering (3.3), (3.4) and Lemmas 3.1–3.3, we obtain

$$\begin{aligned}\frac{1}{4}|\!|\!|G|\!|\!|_\omega^2 \leqslant& a_{MSD}(\omega^{-1}G,G) = a_{MSD}(E,G) + (\omega^{-1}G)(\boldsymbol{x}^*)\\ \leqslant& \frac{1}{8}|\!|\!|G|\!|\!|_\omega^2 + \begin{cases} CN^2\sigma_\beta & \text{if } \boldsymbol{x}^* \in \Omega_s \\ CN\ln N & \text{if } \boldsymbol{x}^* \in \Omega_x \cup \Omega_y \end{cases}.\end{aligned}$$

Thus we are done. □

From (3.5), (3.19) and (3.24) we have

$$\sigma_\beta \geqslant kN^{-1}\ln N, \quad \sigma_\eta \geqslant k\max\{\hat{\varepsilon}_M^{1/2}, N^{-3/4}, \sigma_\eta^*, N^{-1}\ln N, \varepsilon^{1/4}N^{-1/2}\ln^{1/2}N\}$$

where $\sigma_\eta^* = \begin{cases} kN^{-1/2} & \text{if } \hat{\varepsilon}_s \leqslant N^{-2} \\ k\hat{\varepsilon}_s^{-1/2}N^{-3/2} & \text{if } \hat{\varepsilon}_s \geqslant N^{-2} \end{cases}$. Note that if $N \geqslant 4$, then $N^{1/2} \geqslant \ln N$ and

$$N^{-1/2} \geqslant N^{-1}\ln N, \quad N^{-1/2} \geqslant \varepsilon^{1/4}N^{-1/2}\ln^{1/2}N \quad \text{for } N \geqslant 4.$$

Now, we consider the case of $\hat{\varepsilon} = \varepsilon$. Clearly, $\hat{\varepsilon}_M = \hat{\varepsilon}_s = \varepsilon$.

- If $\varepsilon \leqslant N^{-2}$, then $\sigma_\eta^* = kN^{-1/2}$ and we have

$$\begin{aligned}&N^{-1/2} \geqslant \varepsilon^{1/2}, \quad N^{-1/2} \geqslant N^{-3/4},\\ &N^{-1/2} \geqslant N^{-1}\ln N \quad \text{for } N \geqslant 4,\\ &N^{-1/2} \geqslant \varepsilon^{1/4}N^{-1/2}\ln^{1/2}N \quad \text{for } N \geqslant 4.\end{aligned}$$

- If $N^{-2} < \varepsilon \leqslant N^{-3/2}$, then $\sigma_\eta^* = k\varepsilon^{-1/2}N^{-3/2}$ and we have

  $$\varepsilon^{-1/2}N^{-3/2} \geqslant \varepsilon^{1/2}, \quad \varepsilon^{-1/2}N^{-3/2} \geqslant N^{-3/4}.$$

  It means that $\sigma_\eta \geqslant k\max\{\varepsilon^{-1/2}N^{-3/2}, N^{-1}\ln N, \varepsilon^{1/4}N^{-1/2}\ln^{1/2}N\}$. Note that if $N^{-2} < \varepsilon \leqslant N^{-3/2}$,

  $$\begin{aligned} &N^{-1/2} > \varepsilon^{-1/2}N^{-3/2} \\ &N^{-1/2} > N^{-1}\ln N \quad \text{for } N \geqslant 4, \\ &N^{-1/2} > \varepsilon^{1/4}N^{-1/2}\ln^{1/2}N \quad \text{for } N \geqslant 4. \end{aligned}$$

- If $N^{-3/2} < \varepsilon \leqslant N^{-1}$, then $\sigma_\eta^* = k\varepsilon^{-1/2}N^{-3/2}$ and we have

  $$\varepsilon^{1/2} \geqslant \varepsilon^{-1/2}N^{-3/2}, \quad \varepsilon^{1/2} \geqslant N^{-3/4}.$$

  It means that $\sigma_\eta \geqslant k\max\{\varepsilon^{1/2}, N^{-1}\ln N, \varepsilon^{1/4}N^{-1/2}\ln^{1/2}N\}$. Note that if $N^{-3/2} < \varepsilon \leqslant N^{-1}$,

  $$\begin{aligned} &N^{-1/2} > \varepsilon^{1/2} \\ &N^{-1/2} > N^{-1}\ln N \quad \text{for } N \geqslant 4, \\ &N^{-1/2} > \varepsilon^{1/4}N^{-1/2}\ln^{1/2}N \quad \text{for } N \geqslant 4. \end{aligned}$$

These mean that *in the case of $\hat{\varepsilon} = \varepsilon$ we can set $\sigma_\eta \geqslant kN^{-1/2}$.*

If $\hat{\varepsilon} = \begin{cases} \tilde{\varepsilon} & \text{if } \boldsymbol{x} \in \Omega_s, \\ \varepsilon & \text{otherwise,} \end{cases}$ where $\tilde{\varepsilon} := \max\left(\varepsilon, N^{-3/2}\right)$. Clearly, $\hat{\varepsilon}_M = \hat{\varepsilon}_s = \tilde{\varepsilon}$. Note that $\sigma_\eta^* = k\tilde{\varepsilon}^{-1/2}N^{-3/2}$ for $\hat{\varepsilon}_s \geqslant N^{-3/2}$.

- If $\varepsilon \leqslant N^{-3/2}$, then $\hat{\varepsilon}_M = \tilde{\varepsilon} = N^{-3/2}$ and we have

  $$\hat{\varepsilon}_M^{1/2} = \tilde{\varepsilon}^{1/2} = N^{-3/4}, \quad \tilde{\varepsilon}^{-1/2}N^{-3/2} = N^{-3/4}.$$

  It means that $\sigma_\eta \geqslant k\max\{\tilde{\varepsilon}^{1/2}, N^{-1}\ln N, \varepsilon^{1/4}N^{-1/2}\ln^{1/2}N\}$. Note that

  $$\begin{aligned} &\tilde{\varepsilon}^{1/2}\ln^{1/2}N > N^{-1}\ln N \quad \text{for } N \geqslant 4, \\ &\tilde{\varepsilon}^{1/2}\ln^{1/2}N > \varepsilon^{1/4}N^{-1/2}\ln^{1/2}N \quad \text{for } N \geqslant 4. \end{aligned}$$

- If $N^{-3/2} < \varepsilon \leqslant N^{-1}$, then $\tilde{\varepsilon} = \varepsilon > N^{-3/2}$ and we have

  $$\begin{aligned} &\tilde{\varepsilon}^{1/2} > \varepsilon^{-1/2}N^{-3/2}, \\ &\tilde{\varepsilon}^{1/2}\ln^{1/2}N \geqslant N^{-1}\ln N \quad \text{for } N \geqslant 4 \\ &\tilde{\varepsilon}^{1/2}\ln^{1/2}N \geqslant \varepsilon^{1/4}N^{-1/2}\ln^{1/2}N \quad \text{for } N \geqslant 4 \end{aligned}$$

These mean that *we can take* $\sigma_\eta \geqslant k\tilde{\varepsilon}^{1/2}\ln^{1/2} N$.

From the above derivations, we have the following remark.

Remark 3.1. *If we set* $\hat{\varepsilon} = \varepsilon$*, we can take*

$$\sigma_\beta = kN^{-1}\ln N, \quad \sigma_\eta = kN^{-1/2}.$$

If we set $\hat{\varepsilon}$ *as in [3, pg 463], i.e.,*

$$\hat{\varepsilon} = \begin{cases} \tilde{\varepsilon} & \text{if } \boldsymbol{x} \in \Omega_s, \\ \varepsilon & \text{otherwise}, \end{cases}$$

where $\tilde{\varepsilon} := \max\left(\varepsilon, N^{-3/2}\right)$*, we can define*

$$\sigma_\beta = kN^{-1}\ln N, \quad \sigma_\eta = k\tilde{\varepsilon}^{1/2}\ln^{1/2} N.$$

It seems that the definition of σ_β *in [3] is not proper.*

References

[1] C. Johnson, A. H. Schatz, and L. B. Wahlbin. Crosswind smear and pointwise errors in streamline diffusion finite element methods. *Math. Comp.*, 49(179):25–38, 1987.

[2] T. Linß. Layer-adapted meshes for convection–diffusion problems. *Comput. Methods Appl. Mech. Engrg.*, 192(9):1061–1105, 2003.

[3] T. Linß and M. Stynes. The SDFEM on Shishkin meshes for linear convection–diffusion problems. *Numer. Math.*, 87(3):457–484, 2001.

[4] K. Niijima. Pointwise error estimates for a streamline diffusion finite element scheme. *Numer. Math.*, 56(7):707–719, 1990.

[5] H.-G. Roos. Layer-adapted grids for singular perturbation problems. *ZAMM Z. Angew. Math. Me.*, 78(5):291–309, 1998.

[6] H.-G. Roos, M. Stynes, and L. Tobiska. *Numerical Methods for Singularly Perturbed Differential Equations.* Springer-Verlag, Berlin and Heidelberg and New York, 1996.

[7] H.-G. Roos, M. Stynes, and L. Tobiska. *Robust Numerical Methods for Singularly Perturbed Differential Equations*, volume 24 of *Springer Series in Computational Mathematics.* Springer-Verlag, Berlin, 2 edition, 2008.

[8] M. Stynes and L. Tobiska. The SDFEM for a convection–diffusion problem with a boundary layer: optimal error analysis and enhancement of accuracy. *SIAM J. Numer. Anal.*, 41(5):1620–1642, 2003.

[9] Z. M. Zhang. Finite element superconvergence on Shishkin mesh for 2-D convection–diffusion problems. *Math. Comp.*, 72(243):1147–1177, 2003.